

**Amendments to the Claims**

This listing of claims will replace all prior versions and listings of claims in the application:

**Listing of Claims:**

Claims 1-92 (Cancelled)

93 (Currently amended). \* An isolated monoclonal antibody which recognizes IGIF or IL-18 having the following physiochemical properties of (1) to (4):

- (1) Molecular weight  
19,000±5,000 daltons on gel filtration and sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE);
- (2) Isoelectric point (pI)  
4.8±1.0 on chromatofocusing;
- (3) Biological activity  
Inducing the interferon- $\gamma$  production by immunocompetent cells; and
- (4) Amino acid sequence  
Comprising the amino acid sequence of SEQ ID NO:2, wherein Xaa is Met or Thr.

Claim 94 (Cancelled).

95 (Currently amended). A An isolated monoclonal antibody according to claim 93, wherein said IGIF or IL-18 is obtainable from a mammal.

Claims 96 and 97 (Cancelled).

98 (Currently amended). A An isolated monoclonal antibody according to claim 93 or 95 which is an IgG or IgM class antibody.

99 (Currently amended). An isolated antibody according to claim 93 or 95 which is labeled with a radiolabel, an enzyme, or a fluorophore.

100 (Currently amended). An isolated antibody according to claim 93 or 95 which is capable of inhibiting the biological activity of IGIF or IL-18.

101 (Previously presented). A hybridoma which produces a monoclonal antibody according to claim 93 or 95.

102 (Previously presented). A method for producing a monoclonal antibody which comprises culturing a hybridoma according to claim 101 *in vitro* or *in vivo* under conditions suitable to promote production of the antibody and recovering the antibody so produced.

103 (Previously presented). A method according to claim 102, further comprising the step of subjecting the antibody to one or more processes selected from the group consisting of salting out, dialysis, filtration, concentration, centrifugation, separatory sedimentation,

gel filtration chromatography, ion- exchange chromatography, HPLC, affinity chromatography, gel electrophoresis, and isoelectric focusing.

104(Previously amended). A method for determining the presence of IGIF or IL-18 in a sample, comprising the steps of:

contacting a sample suspected to contain IGIF or IL-18 with an antibody according to claim 93 or 95 under conditions suitable to promote the specific binding of the antibody to IGIF or IL-18 to form an immune complex; and

detecting any such immune complex which is so formed.

105(Previously presented). A method according to claim 104, wherein the antibody is immobilized on an insoluble matrix or substrate.

106(Previously presented). A method according to claim 104, wherein the antibody is labeled with a radiolabel, an enzyme, or a fluorophore.

107(Previously presented). A method according to claim 104, further comprising the step of quantifying the amount of IGIF or IL-18 present in the sample.

108(Previously presented). A method according to claim 104, wherein the IGIF or IL-18 has the amino acid sequence shown in SEQ ID NO:2, wherein Xaa is Met or Thr.

109(Previously presented). A method for purifying IGIF or IL-18 from a sample containing other components, comprising the steps of:

contacting the sample with a monoclonal antibody according to claim 93 or 95 under conditions suitable to promote the specific binding of the antibody to IGIF or IL-18 to form an immune complex; and separating the immune complex from at least one of the other components in the sample.

110(Previously presented). A method according to claim 109, further comprising the step of recovering the IGIF or IL-18 from the immune complex.

111(Previously presented). A method according to claim 109, wherein the antibody is immobilized on an insoluble matrix.

112(Previously presented). A method according to claim 109, wherein the contacting step is effected by applying the sample to a chromatography column comprising an insoluble matrix.

113(Previously presented). A method according to claim 112, further comprising the step of recovering the IGIF or IL-18 from the chromatography column.

114(Previously presented). A method according to claim 113, wherein the IGIF or IL-18 is recovered in nearly quantitative yield and with a purity of at least 95%.

115(Previously presented). A method according to claim 109, wherein the IGIF or IL-18 has the amino acid sequence shown in SEQ ID NO:2, wherein Xaa is Met or Thr.

116(Previously presented). A method of inhibiting the biological activity of IGIF or IL-18, comprising the step of contacting an antibody according to claim 100, with the IGIF or IL-18.

117(Previously presented). A method according to claim 116, wherein the IGIF or IL-18 has the amino acid sequence shown in SEQ ID NO:2, wherein Xaa is Met or Thr.

118(Currently amended). ~~A~~ An isolated monoclonal antibody, which recognizes ~~specific to~~ interferon-gamma (IFN- $\gamma$ ) inducing protein, also known as IGIF and IL-18, having the amino acid sequence shown in SEQ ID NO:2, wherein Xaa is Met or Thr.

119(Currently amended). ~~A~~ An isolated monoclonal antibody according to claim 95, wherein said mammal is mouse.

120(Currently amended). An isolated antibody obtainable by using, as an antigen, IGIF or IL-18 having the following physiochemical properties of (1) to (4):

- (1) Molecular weight  
19,000 $\pm$ 5,000 daltons on gel filtration and sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE);
- (2) Isoelectric point (pI)

Appln. No. 09/050,249  
Amd. dated October 26, 2006  
Reply to Office Action of July 26, 2006

4.8±1.0 on chromatofocusing;

(3) Biological activity

Inducing interferon- $\gamma$  production by immunocompetent  
cells; and

(4) Amino acid sequence

Comprising the amino acid sequence of SEQ ID NO:2, wherein Xaa is Met  
or Thr.